

THE CONTINUITY OF THE INVERSION AND THE STRUCTURE OF MAXIMAL SUBGROUPS IN COUNTABLY COMPACT TOPOLOGICAL SEMIGROUPS

OLEG GUTIK, DUŠAN PAGON, AND DUŠAN REPOVŠ

ABSTRACT. In this paper we search for conditions on a countably compact (pseudo-compact) topological semigroup under which: (i) each maximal subgroup $H(e)$ in S is a (closed) topological subgroup in S ; (ii) the Clifford part $H(S)$ (i.e. the union of all maximal subgroups) of the semigroup S is a closed subset in S ; (iii) the inversion $\text{inv}: H(S) \rightarrow H(S)$ is continuous; and (iv) the projection $\pi: H(S) \rightarrow E(S)$, $\pi: x \mapsto xx^{-1}$, onto the subset of idempotents $E(S)$ of S , is continuous.

In this paper all topological spaces will be assumed to be Hausdorff. We shall follow the terminology of [7, 8, 11]. We shall denote the cardinality of continuum by $\mathfrak{c}$ and the topological closure of subset A in a topological space by $\overline{A}$. We shall call a T_3 -space a regular topological space.

A topological space X is said to be *countably compact* if any countable open cover of X contains a finite subcover [11]. A topological space X is called *pseudocompact* if each continuous real-valued function on X is bounded [11]. A topological space X is said to be *sequential* if each non-closed subset A of X contains a sequence of points $\{x_n\}_{n=1}^{\infty}$ that converges to some point $x \in X \setminus A$. Obviously, a topological space X is sequential if a subset A of X is closed if and only if together with any convergent sequence A contains its limit [11]. A topological space X is called *sequentially compact* if each sequence $\{x_n\}_{n=1}^{\infty} \subset X$ has a convergent subsequence [11].

We recall that the Stone-Čech compactification of a Tychonoff space X is a compact Hausdorff space βX containing X as a dense subspace so that each continuous map $f: X \rightarrow Y$ to a compact Hausdorff space Y extends to a continuous map $\tilde{f}: \beta X \rightarrow Y$ [11].

A *semigroup* is a set with a binary associative operation. An element e of a semigroup S is called an *idempotent* if $e \circ e = e$. If S is a semigroup, then we denote the subset of all idempotents of S by $E(S)$. A semigroup S is called *inverse* if for any $x \in S$ there exists a unique $y \in S$ such that $xyx = x$ and $xyx = y$. Such an element y is called *inverse* of x and is denoted by x^{-1} . If S is an inverse semigroup, then the map which takes $x \in S$ to the inverse element of x is called the *inversion* and will be denoted by inv .

If S is a semigroup and e is an idempotent in S , then e lies in the maximal subgroup $H(e)$ with the identity e . If a semigroup S is a union of groups then S is called *Clifford*. On a Clifford semigroup $S = \bigcup \{H(e) \mid e \in E(S)\}$ the inversion $\text{inv}: S \rightarrow S$ is defined which maps each element $x \in H(e)$ to its inverse element x^{-1} in $H(e)$. We also observe that on any Clifford semigroup the *projection* $\pi: S \rightarrow E(S)$, $\pi(x) = x \cdot x^{-1}$, is defined. For a semigroup S let

$$\begin{aligned} H(S) &= \bigcup_{e \in E(S)} \{H(e) \mid H(e) \text{ is a maximal subgroup in } S \text{ with identity } e\} = \\ &= \{s \in S \mid \text{there exists } y \in S \text{ such that } xy = yx, xyx = x, yxy = y\}. \end{aligned}$$

A topological space S which is algebraically a semigroup with a continuous semigroup operation is called a *topological semigroup*. A *topological inverse semigroup* is a topological semigroup S that is algebraically an inverse semigroup with continuous inversion. If τ is a topology on a (inverse) semigroup S such that (S, τ) is a topological (inverse) semigroup, then τ is called a *(inverse) semigroup topology* on S . By a *paratopological group* we understand a pair (G, τ) consisting of a group G and a topology τ on G making the group operation on G continuous. A paratopological group G with continuous inversion is called a *topological group*.

Date: July 22, 2009.

2000 Mathematics Subject Classification. Primary 22A15, 54H10, 54D55. Secondary 22A05, 54D30, 54D40.

Key words and phrases. Topological semigroup, topological inverse semigroup, sequential space, sequentially compact space, countably compact space, pseudocompact space, regular space, quasi-regular space, subgroup, closure, inversion, paratopological group, topological group, Clifford semigroup, topologically periodic semigroup, $\text{MA}_{\text{countable}}$, selective ultrafilter.

Finite semigroups and compact topological semigroups have similar properties. For example every finite semigroup and every compact topological semigroup contain idempotents and minimal ideals [26], which are completely simple semigroups [25, 27], and every (0-)simple compact topological (and hence finite) semigroup is completely (0-)simple [20, 25, 27]. Also, a cancellative compact topological (and hence finite) semigroup is a topological group [19].

Compact topological semigroups do not contain the bicyclic semigroup [15]. Gutik and Repovš [14] proved that a countably compact topological inverse semigroup does not contain the bicyclic semigroup. Banakh, Dimitrova and Gutik [2] showed that no topological semigroup S with countably compact square $S \times S$ contains the bicyclic semigroup. They also constructed in [3] a consistent example a countably compact topological semigroup S which contains the bicyclic semigroup.

It is well known that the closure of a (commutative) subsemigroup of a topological semigroup is a (commutative) subsemigroup [7, Vol. 1, P. 9]. Note that the closure of a subgroup in a topological semigroup is not necessarily a subgroup. But in the case when S is a compact topological semigroup or a topological inverse semigroup, the closure of a subgroup in S is a subgroup, moreover every maximal subgroup of S is closed (see [7, Vol. 1, Theorem 1.11] and [9]).

Also, every compact subgroup of a topological semigroup with induced topology is a topological semigroup. The results when the inversion is continuous in a topological semigroup which is algebraically a group (i. e. a paratopological group) have been extended to some classes of “compact-like” paratopological groups, in particular: regular locally compact paratopological groups [10], regular countably compact paratopological groups [22], quasi-regular pseudo-compact paratopological groups [22], topologically periodic Hausdorff countably compact paratopological groups [5], Čech-complete paratopological groups [6], strongly Baire semi-topological groups [16].

On the other hand, Ravsky [21], using a result of Koszmider, Tomita and Watson [17], constructed a MA-example of a Hausdorff countably compact paratopological group failing to be a topological group. Also Grant [13] and Yur’eva [29] showed that a Hausdorff cancellative sequentially countably compact topological semigroup is a topological group. Bokalo and Guran [5] established that an analogous theorem is true for cancellative sequentially compact semigroups. Robbie and Svetlichny [23] constructed a CH-example of a countably compact topological semigroup which is not a topological group.

In summary (see [7]), for a compact topological semigroup S the following conditions hold:

- (1) each maximal subgroup $H(e)$ in S is a compact topological subgroup in S ;
- (2) the subset $H(S)$ is closed in S ;
- (3) the inversion map $\text{inv}: H(S) \rightarrow H(S)$ is continuous; and
- (4) the projection $\pi: H(S) \rightarrow E(S)$ is continuous.

Since sequential compactness, countable compactness and pseudo-compactness are generalization of compactness, it is natural to pose the following question: *For which compact-like topological semigroups do the conditions (1)–(4) above hold?*

In this paper we shall answer this question by giving sufficient conditions on a countably compact (pseudo-compact) topological semigroup under which: (i) each maximal subgroup $H(e)$ in S is a (closed) topological subgroup in S ; (ii) the Clifford part $H(S)$ of the semigroup S is a closed subset in S ; and (iii) the inversion $\text{inv}: H(S) \rightarrow H(S)$ and the projection $\pi: H(S) \rightarrow E(S)$ are continuous.

A topological group G is called *totally bounded* if for any open neighbourhood U of the identity e of G there exists a finite subset A in G such that $A \cdot U = G$ (see [28]).

Theorem 1. *Let S be a Tychonoff topological semigroup with the pseudocompact square $S \times S$. Then S embeds into a compact topological semigroup and the following conditions hold:*

- (i) *the inversion $\text{inv}: H(S) \rightarrow H(S)$ is continuous;*
- (ii) *the projection $\pi: H(S) \rightarrow E(S)$ is continuous; and*
- (iii) *for each idempotent $e \in E(S)$ the maximal subgroup $H(e)$ is a totally bounded topological group.*

Proof. By Theorem 1.3 from [1], for any topological semigroup S with the pseudocompact square $S \times S$ the semigroup operation $\mu: S \times S \rightarrow S$ extends to a continuous semigroup operation $\beta\mu: \beta S \times \beta S \rightarrow \beta S$, so S is a subsemigroup of the compact topological semigroup βS .

(i) Let

$$\text{Gr}_{\text{inv}}(H(\beta S)) = \{(x, y) \in S \times S \mid y = x^{-1}\}$$

be the graph of the inversion in $H(\beta S)$. Since βS is a topological semigroup and

$$\text{Gr}_{\text{inv}}(H(\beta S)) = \{(x, y) \in S \times S \mid xyx = x, yxy = y \text{ and } xy = yx\},$$

the graph $\text{Gr}_{\text{inv}}(H(\beta S))$ is a compact subset of $\beta S \times \beta S$.

Consider the natural projections $\text{pr}_1: \beta S \times \beta S \rightarrow \beta S$ and $\text{pr}_2: \beta S \times \beta S \rightarrow \beta S$ onto the first and the second coordinates, respectively. It follows from the compactness of $\text{Gr}_{\text{inv}}(H(\beta S))$ that $\text{pr}_1: \text{Gr}_{\text{inv}}(H(\beta S)) \rightarrow H(\beta S)$ and $\text{pr}_2: \text{Gr}_{\text{inv}}(H(\beta S)) \rightarrow H(\beta S)$ are homeomorphisms. Consequently, the inversion $\text{inv}|_{H(\beta S)} = \text{pr}_2 \circ (\text{pr}_1)^{-1}: H(\beta S) \rightarrow H(\beta S)$ is continuous, being a composition of homeomorphisms. Therefore the inversion $\text{inv}: H(S) \rightarrow H(S)$ is continuous as a restriction of a continuous map.

(ii) The projection $\pi: H(S) \rightarrow E(S)$ is continuous as a composition of two continuous maps.

(iii) Given an idempotent $e \in E(S)$, consider the maximal subgroup $H_\beta(e)$ in βS containing e . Then by Theorems 1.11 and 1.13 from [7, Vol. 1], $H_\beta(e)$ is a compact topological group and since $H(e)$ is a subgroup of $H_\beta(e)$ the inversion $\text{inv}: H(e) \rightarrow H(e)$ is continuous, and $H(e)$ is a totally bounded topological group, see [28]. $\square$

Theorem 1 implies the following:

Corollary 2. *If S is a Tychonoff Clifford topological semigroup with the pseudocompact square $S \times S$ then the inversion in S is continuous.*

An element x of a topological semigroup S is called *topologically periodic* if for any open neighbourhood $U(x)$ of x there exists an integer $n \geq 2$ such that $x^n \in U(x)$. A topological semigroup S is called *topologically periodic* if any element of S is topologically periodic.

Remark 3. The following observation implies that *an element of any (not necessarily Hausdorff) topological semigroup S is topologically periodic if and only if for any integer $n \geq 2$ and for any open neighbourhood $U(x)$ of x there exists an integer $m \geq n$ such that $x^m \in U(x)$* . Let $k \geq 2$ be an integer such that $x^k \in U(x)$. Then the continuity of the semigroup operation implies that there exists an open neighbourhood $V(x)$ of x such that $(V(x))^k \subseteq U(x)$. Since x is topologically periodic there exists an integer $m \geq 2$ such that $x^m \in V(x)$. Hence we have $x^{km} \in (V(x))^k \subseteq U(x)$ and $km \geq 4 = 2^2$. Proceeding by induction, we can find an integer $p \geq 2^n > n$ such that $x^p \in U(x)$.

Theorem 4. *Let S be a Hausdorff topological semigroup with the countably compact square $S \times S$. Then:*

- (i) *each maximal subgroup $H(e)$ of S is a countably compact topological group; and*
- (ii) *the subset $H(S)$ is countably compact.*

Proof. (i) Let $H(e)$ be any maximal subgroup of S . Since the semigroup operation in S is continuous the subset

$$G = \{(x, y) \in S \times S \mid xy = yx = e, xe = ex = x, ye = ey = y\}$$

is closed in $S \times S$ and Theorem 3.10.4 from [11] implies that G is a countably compact subset in $S \times S$. Consider the natural projection $\text{pr}_1: S \times S \rightarrow S$ onto the first coordinate. Since $\text{pr}_1(G) = H(e)$ and the projection $\text{pr}_1: S \times S \rightarrow S$ is a continuous map Theorem 3.10.5 from [11] implies that $H(e)$ is a countably compact subspace of S .

Next, we show that $H(e)$ is a topologically periodic paratopological group. Let x be an arbitrary element of the subgroup $H(e)$ and $U(x)$ be any open neighbourhood of x . We consider the sequence $\{(x^{n+1}, x^{-n})\}_{n=1}^\infty$ in $H(e) \times H(e) \subseteq S \times S$. The countable compactness of $S \times S$ guarantees that this sequence has an accumulation point $(a, b) \in S \times S$. Since $x^{n+1} \cdot x^{-n} = x$, the continuity of the semigroup operation on S guarantees that $ab = x$. Then for any open neighbourhood $U(x)$ of x in S there exist open neighbourhoods $U(a)$ and $U(b)$ of the point a and b in S , respectively, such that $U(a)U(b) \subseteq U(x)$. Since (a, b) is an accumulation point of the sequence $\{(x^{n+1}, x^{-n})\}_{n=1}^\infty$ in $S \times S$, there exist positive integers m and n such that $x^m \in U(a)$, $x^{-n} \in U(b)$ and $m \geq n + 2$. Hence we get that $x^m \cdot x^{-n} = x^{m-n} \in U(a) \cdot U(b) \subseteq U(x)$ and $m - n \geq 2$. Therefore $H(e)$ is a topologically periodic paratopological group. By Bokalo-Guran Theorem (see [5, Theorem 3]) any countably compact paratopological group is a topological group. Consequently, $H(e)$ is a countably compact topological group.

(ii) Since the semigroup operation in S is continuous,

$$H = \{(x, y) \in S \times S \mid xyx = x, yxy = y, xy = yx \in E(S)\}$$

is a closed subset in $S \times S$ and Theorem 3.10.4 from [11] implies that H is a countably compact subset in $S \times S$. Consider the natural projection $\text{pr}_1: S \times S \rightarrow S$ onto the first coordinate. Since $\text{pr}_1(H) = H(S)$ and the projection $\text{pr}_1: S \times S \rightarrow S$ is a continuous map, Theorem 3.10.5 from [11] implies that $H(S)$ is a countably compact subspace of S . $\square$

Proposition 5. *Let x be a topologically periodic element of a maximal subgroup $H(e)$ with the unity e in a topological semigroup S . Then the inversion $\text{inv}: H(S) \rightarrow H(S)$ is continuous at x if and only if it is continuous at the idempotent e .*

Proof. We follow the argument of [4]. Let $U(x^{-1})$ be any open neighbourhood of the inverse element x^{-1} of x in S . Since the semigroup operation in S is continuous there exist open neighbourhoods $V(x^{-1})$ and $V(e)$ of x^{-1} and e in $H(S)$, respectively, such that $V(x^{-1}) \cdot V(e) \subseteq U(x^{-1})$. Since the inversion is continuous at idempotent e there exists an open neighbourhood $W(e)$ of e in $H(S)$ such that $(W(e))^{-1} \subseteq V(e)$.

Also, the continuity of the semigroup operation implies that there exists an open neighbourhood $N(x)$ of x in $H(S)$ such that $x^{-1} \cdot N(x) \cdot x^{-1} \subseteq V(x^{-1})$ and $N(x) \cdot x^{-1} \subseteq W(e)$. The topological periodicity of x implies that there exists a positive integer n such that $x^{n+2} \in N(x)$. Then we have that

$$x^{n+1} = x^{n+2} \cdot x^{-1} \in N(x) \cdot x^{-1} \subseteq W(e)$$

and

$$x^n = x^{-1} \cdot x^{n+2} \cdot x^{-1} \in x^{-1} \cdot N(x) \cdot x^{-1} \subseteq V(x^{-1}).$$

Since S is a topological semigroup there exists an open neighbourhood $P(x)$ of x in S such that $(H(S) \cap P(x))^{n+1} \subseteq W(e)$ and $(H(S) \cap P(x))^n \subseteq V(x^{-1})$. Therefore we get

$$\begin{aligned} (H(S) \cap P(x))^{-1} &\subseteq (H(S) \cap P(x))^n \cdot \left((H(S) \cap P(x))^{n+1} \right)^{-1} \subseteq V(x^{-1}) \cdot (W(e))^{-1} \subseteq \\ &\subseteq V(x^{-1}) \cdot V(e) \subseteq U(x^{-1}), \end{aligned}$$

and hence the inversion is continuous at the point x . $\square$

Proposition 5 implies the following:

Corollary 6. *The inversion in a topologically periodic Clifford topological semigroup S is continuous if and only if it is continuous at any idempotent of the semigroup S .*

Theorem 7. *Let S be a regular topological semigroup with the countably compact square $S \times S$. Then:*

- (i) *the inversion $\text{inv}: H(S) \rightarrow H(S)$ is continuous; and*
- (ii) *the projection $\pi: H(S) \rightarrow E(S)$ is continuous.*

Proof. (i) By Proposition 5 it is sufficient to show that the inversion $\text{inv}: H(S) \rightarrow H(S)$ is continuous at any point of the set $E(S)$.

Fix any $e \in E(S)$. Let $U(e)$ be any open neighbourhood of e in S . Since the topological space of the semigroup S is regular, the continuity of the semigroup operation of S implies that there exists a sequence of open neighbourhoods $\{U_i(e)\}_{i=1}^{\infty}$ of the idempotent e in S such that $\overline{U_1(e)} \subseteq U(e)$ and $\overline{(U_n(e))^m} \subseteq U_{n-1}(e)$ for any positive integer n and all $m = 1, \dots, n$. Let $F = \bigcap_{n=1}^{\infty} \overline{U_n(e)}$.

We shall show that $(F \cap H(S))^{-1} \subseteq F$. Let x be any element of the set $F \cap H(S)$. Since the set F is closed to prove that $x^{-1} \in F$ it is sufficient to show that $V(x^{-1}) \cap F \neq \emptyset$ for any open neighbourhood $V(x^{-1})$ of the point x^{-1} . The continuity of the semigroup operation in S and the equality $x^{-1} = x^{-1} \cdot x \cdot x^{-1}$ imply that there exists an open neighbourhood $V(x)$ of the point x in S such that $x^{-1} \cdot V(x) \cdot x^{-1} \subseteq V(x^{-1})$. By Theorem 4 (i) the element x of S is topologically periodic, and hence there exists a positive integer $n \geq 2$ such that $x^n \in V(x)$. Then we have

$$x^{n-2} = x^{-1} \cdot x^n \cdot x^{-1} \in x^{-1} \cdot V(x) \cdot x^{-1} \subseteq V(x^{-1})$$

and

$$x^{n-2} \in F^{n-2} \subseteq \bigcap_{i=n-2}^{\infty} (U_i(e))^{n-2} \subseteq \bigcap_{i=n-2}^{\infty} U_{i-1}(e) \subseteq F.$$

Hence $V(x^{-1}) \cap F \neq \emptyset$ and since F is a closed subset in S we have that $x^{-1} \in F$. This implies that the inclusion $(F \cap H(S))^{-1} \subseteq F$ holds.

Later we shall show that $(U_n(e) \cap H(S))^{-1} \subseteq U(e) \cap H(S)$ for some positive integer n . Suppose to the contrary that $(U_n(e) \cap H(S))^{-1} \not\subseteq U(e) \cap H(S)$ for any positive integer n . Then there exists a sequence $\{x_n\}_{n=1}^{\infty}$ in $H(S)$ such that $x_n \in U_n(e) \setminus (U(e))^{-1}$ for all positive integers n .

The countable compactness of the square $S \times S$ implies that the sequence $\{(x_n, x_n^{-1})\}_{n=1}^{\infty}$ has a cluster point (a, b) in $S \times S$. The continuity of the semigroup operation in S implies that

$$a \cdot b = b \cdot a = f, \quad a \cdot b \cdot a = a, \quad b \cdot a \cdot b = b,$$

and hence $a, b \in H(f)$ for some idempotent f in S . Therefore $b = a^{-1} \in F^{-1} \cap H(S) \subseteq F$. Then $(a, b) \in F \times F \subseteq U(e) \times U(e)$. Since (a, b) is a cluster point of the sequence $\{(x_n, x_n^{-1})\}_{n=1}^{\infty}$, there exists a positive integer n such that $(x_n, x_n^{-1}) \in U(e) \times U(e)$.

Therefore we have that $x_n \in (U(e))^{-1}$ which contradicts the choice of the sequence $\{x_n\}_{n=1}^{\infty}$. The obtained contradiction implies that $(U_n(e) \cap H(S))^{-1} \subseteq U(e) \cap H(S)$ for some positive integer n , and hence the inversion $\text{inv}: H(S) \rightarrow H(S)$ is continuous.

(ii) The projection $\pi: H(S) \rightarrow E(S)$ is continuous as a composition of two continuous maps. $\square$

Theorem 7 implies the following corollary generalizing a result of [4].

Corollary 8. *The inversion in a regular Clifford topological semigroup with the countably compact square is continuous.*

Let S be a topological semigroup and $e \in E(S)$. We shall say that the semigroup S is *inversely regular at e* if for any open neighbourhood $U(e)$ of e there exists an open neighbourhood $W(e)$ of e such that $\overline{(W(e) \cap H(S))^{-1}} \subseteq (U(e) \cap H(S))^{-1}$. A topological semigroup S with non-empty subsets of idempotents is called *inversely regular* if it is inversely regular at each idempotent of S [4].

Theorem 9. *Let S be a topologically periodic Hausdorff topological semigroup. If S is inversely regular and countably compact, then:*

- (i) *the inversion $\text{inv}: H(S) \rightarrow H(S)$ is continuous;*
- (ii) *the projection $\pi: H(S) \rightarrow E(S)$ is continuous.*

Proof. (i) Fix any idempotent e in S . Let $U(e)$ be any open neighbourhood of e in S . Since the semigroup operation in S is continuous and S is inversely regular we construct inductively two sequences $\{U_n(e)\}_{n=1}^{\infty}$ and $\{W_n(e)\}_{n=1}^{\infty}$ of open neighbourhoods of the idempotent e such that $(U_n(e))^i \subseteq W_{n-1}(e)$ and $\overline{(W_n(e) \cap H(S))^{-1}} \subseteq (U_n(e) \cap H(S))^{-1}$ for all positive integers n and $i = 1, 2, \dots, n$.

Let $F = \bigcap_{n=1}^{\infty} \overline{(W_n(e) \cap H(S))^{-1}}$. Then we have that

$$F = \bigcap_{n=1}^{\infty} \overline{(W_n(e) \cap H(S))^{-1}} \subseteq \bigcap_{n=2}^{\infty} (U_n(e) \cap H(S))^{-1} \subseteq \bigcap_{n=2}^{\infty} (W_{n-1}(e) \cap H(S))^{-1} \subseteq F.$$

We shall show that $F^{-1} = F$. Let x be arbitrary element of F . Since the set F is closed it sufficient to prove that $V(x^{-1}) \cap F \neq \emptyset$ for any open neighbourhood $V(x^{-1})$ of the point x^{-1} . Since the semigroup S is topologically periodic there exists a positive integer $n \geq 2$ such that $x^{n-2} \in V(x^{-1})$ (see proof of Theorem 7). Then we have

$$\begin{aligned} x^{n-2} \in F^{n-2} &\subseteq \bigcap_{k=n-2}^{\infty} \left((U_k(e) \cap H(S))^{-1} \right)^{n-2} \subseteq \bigcap_{k=n-2}^{\infty} \left((U_k(e) \cap H(S))^{n-2} \right)^{-1} \subseteq \\ &\subseteq \bigcap_{k=n-2}^{\infty} (W_{k-1}(e) \cap H(S))^{-1} \subseteq F. \end{aligned}$$

Hence $V(x^{-1}) \cap F \neq \emptyset$ and since F is a closed subset in S we have that $x^{-1} \in F$. This implies that the inclusion $F^{-1} \subseteq F$ holds. Then after the inversion we get that $F \subseteq F^{-1}$. Therefore we get that

$$\bigcap_{n=1}^{\infty} (W_n(e) \cap H(S))^{-1} = F = F^{-1} = \bigcap_{n=1}^{\infty} \left(\overline{(W_n(e) \cap H(S))^{-1}} \right)^{-1} \subseteq \bigcap_{n=1}^{\infty} (U_n(e) \cap H(S)) \subseteq U(e).$$

Since the space of the semigroup S is countably compact there exists a positive integer n such that

$$F \subseteq (W_n(e) \cap H(S))^{-1} \subseteq \overline{(W_n(e) \cap H(S))^{-1}} \subseteq U(e).$$

This implies that the inversion is continuous at the idempotent e .

(ii) The projection $\pi: H(S) \rightarrow E(S)$ is continuous as a composition of two continuous maps. $\square$

We recall that a map $f: X \rightarrow Y$ between topological space is called *sequentially continuous* if $\lim_{n \rightarrow \infty} f(x_n) = f(\lim_{n \rightarrow \infty} x_n)$ for any convergent sequence $\{x_n\}_{n=1}^\infty$ in X . Obviously, a composition of two sequentially continuous maps is a continuous map. A subset F of a topological space X is called *sequentially closed* if no sequence in F converges to a point not in F [12].

Theorem 10. *Let S be a Hausdorff countably compact topological semigroup. Then the following conditions hold:*

- (i) *each maximal subgroup $H(e)$, $e \in E(S)$, is sequentially closed in S ;*
- (ii) *the subset $H(S)$ is sequentially closed in S ;*
- (iii) *the inversion $\text{inv}: H(S) \rightarrow H(S)$ is sequentially continuous; and*
- (iv) *the projection $\pi: H(S) \rightarrow E(S)$ is sequentially continuous.*

Proof. (i) Suppose to the contrary that there exists a maximal subgroup $H(e)$ in S which is not sequentially closed subset in S . Then there exists a sequence $\{x_n\}_{n=1}^\infty \subset H(e)$ which converges to $x \notin H(e)$. We put $A = \{x_n\}_{n=1}^\infty \cup \{x\}$. Since the sequence $\{x_n\}_{n=1}^\infty \subset H(e)$ converges to x the set A with the induced from S topology is a compact space. Then by Corollary 3.10.14 from [11], $A \times S$ is a countably compact space.

The countable compactness of $A \times S$ implies that the sequence $\{(x_n, x_n^{-1})\}_{n=1}^\infty$ has a cluster point (a, b) in $A \times S$. The continuity of the semigroup operation in S implies that $ab = e$, $aba = a$, $bab = b$ and hence $a = b^{-1}$ and $a, b \in H(e)$. The Hausdorff property of S implies that $x = a$ and hence $x \in H(e)$. The obtained contradiction implies assertion (i).

(ii) We argue exactly as a previous case. Suppose to the contrary that $H(S)$ is not a sequentially closed subset in S . Then there exists a sequence $\{x_n\}_{n=1}^\infty \subset H(S)$ which converges to $x \notin H(S)$. We put $A = \{x_n\}_{n=1}^\infty \cup \{x\}$. Since the sequence $\{x_n\}_{n=1}^\infty \subset H(S)$ converges to x the set A with the induced from S topology is a compact space. Then by Corollary 3.10.14 from [11], $A \times S$ is a countably compact space.

The countable compactness of $A \times S$ implies that the sequence $\{(x_n, x_n^{-1})\}_{n=1}^\infty$ has a cluster point (a, b) in $A \times S$. The continuity of the semigroup operation in S implies that $ab = e$, $aba = a$, $bab = b$ for some idempotent $e \in E(S)$ and hence $a = b^{-1}$ and $a, b \in H(e)$. The Hausdorff property of S implies that $x = a$ and hence $x \in H(e) \subseteq H(S)$. The obtained contradiction implies assertion (ii).

(iii) The sequential continuity of the inversion $\text{inv}: H(S) \rightarrow H(S)$ will follow as soon as we prove that for any countable compactum $C \subset H(S)$ the restriction $\text{inv}|_C$ is continuous. Let

$$\text{Gr}_{\text{inv}}(S) = \{(x, y) \in S \times S \mid y = x^{-1}\}$$

be the graph of the inversion. Since S is a topological semigroup and

$$\text{Gr}_{\text{inv}}(S) = \{(x, y) \in S \times S \mid xyx = x, yxy = y \text{ and } xy = yx\},$$

the graph $\text{Gr}_{\text{inv}}(S)$ is a closed subset of $S \times S$.

Since C is a metrizable compactum we can apply Corollary 3.10.14 from [11] to conclude that $C \times S$ is a countably compact space. Then the closedness of $\text{Gr}_{\text{inv}}(S)$ in the space $S \times S$ implies that the space $G = (C \times S) \cap \text{Gr}_{\text{inv}}(S)$ is countably compact and being countable, is compact.

Consider the natural projections $\text{pr}_1: S \times S \rightarrow S$ and $\text{pr}_2: S \times S \rightarrow S$ onto the first and the second coordinates, respectively. It follows from the compactness of G that $\text{pr}_1: G \rightarrow C$ and $\text{pr}_2: G \rightarrow C^{-1}$ are homeomorphisms. Consequently, $\text{inv}|_C = \text{pr}_2 \circ (\text{pr}_1)^{-1}: C \rightarrow C^{-1}$ is continuous, being a composition of homeomorphisms.

(iv) The projection $\pi: H(S) \rightarrow E(S)$ is sequentially continuous as a composition of two sequentially continuous maps. $\square$

Theorem 10 implies the following:

Corollary 11. *Let S be a Hausdorff Clifford countably compact topological semigroup. Then the following conditions hold:*

- (i) *each maximal subgroup $H(e)$ is sequentially closed in S ;*
- (ii) *the inversion $\text{inv}: S \rightarrow S$ is sequentially continuous; and*
- (iii) *the projection $\pi: S \rightarrow E(S)$ is sequentially continuous.*

We observe that any sequentially compact (and any hence sequential countably compact) topological semigroup contains an idempotent (see [2, Theorem 8]). For sequential countably compact semigroup Theorem 10 implies the following:

Corollary 12. *Let S be a Hausdorff sequential countably compact topological semigroup. Then the following conditions hold:*

- (i) *each maximal subgroup $H(e)$ is closed in S ;*
- (ii) *the subset $H(S)$ is closed in S ;*
- (iii) *the inversion $\text{inv}: H(S) \rightarrow H(S)$ continuous; and*
- (iv) *the projection $\pi: H(S) \rightarrow E(S)$ is continuous.*

The following example shows that the closure of a subgroup of a countably compact topological semigroup need not be a subgroup.

Example 13. Assume $\text{MA}_{\text{countable}}$ holds. Let $(\mathbb{R}, +)$ be the additive topological group of real number with usual topology and $\mathbb{Z}$ the discrete additive group of integer. Then $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ is a topological group. Let G be the group consisting of all $y \in \mathbb{T}^{\mathfrak{c}}$ such that there exists $\mu \in \mathfrak{c}$ such that $y(\mu)$ is the identity of $\mathbb{T}$ for each $\alpha > \mu$.

There exists $x \in \mathbb{T}^{\mathfrak{c}}$ such that $S = \{nx + y \mid n \in \omega \text{ and } y \in G\}$ is the semigroup with two-sided cancelation but S is not a group, see [23, 24]. Since G is dense subgroup in $\mathbb{T}^{\mathfrak{c}}$, G is dense in S . Tomita [24] showed that the existence of an element x in S is independent of ZFC. Also Madariaga-Garcia and Tomita [18] show that the semigroup S can be constructed from $\mathfrak{c}$ selective ultrafilters.

A topological space X is called *quasi-regular* if for any non-empty open subset U in X there exists a non-empty open set $V \subseteq U$ in X such that $\overline{V} \subseteq U$. The following example shows that there exists a Hausdorff quasi-regular Clifford inverse countably compact topological semigroup S with the discontinuous inversion and discontinuous projection $\pi: S \rightarrow E(S)$.

Example 14 ([4]). Let ω_1 be the smallest uncountable ordinal and $[0, \omega_1)$ be the well-ordered sets of all countable ordinals, endowed with the natural order topology (see [11, Example 3.10.16]). It is well-known that $[0, \omega_1)$ is a sequentially compact topological space (see: [11], Example 3.10.16) and simple verification shows that the semilattice operation $\min$ is continuous on $[0, \omega_1)$.

Let $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$ be the unit circle in the complex plane with the usual topology and let $\mathbb{T}$ endowed with the operation of multiplication of complex numbers. Then by Theorem 3.10.35 from [11] the product $A = [0, \omega_1) \times \mathbb{T}$ is an Hausdorff sequentially compact commutative topological inverse semigroup, as the Cartesian product of a topological semilattice and a commutative topological group.

Let $x = (\omega_1, 1) \notin A$. Put $S = A \cup \{x\}$ and define a topology τ on S letting A be a subspace of S and $U \subset S$ be a neighborhood of x if there are a positive real $\varepsilon > 0$ and a countable ordinal α such that $U \supset U(\alpha, \varepsilon)$ where

$$U(\alpha, \varepsilon) = \{x\} \cup \{(\beta, e^{i\varphi}) \mid \alpha < \beta < \omega_1, 0 < \varphi < \varepsilon\}.$$

Extend the semigroup operation to S letting $x \cdot x = x$ and $x \cdot a = a \cdot x = a$ for all $a \in A$. It is easy to see that S is a sequentially compact space and the semigroup operation "·" is continuous and commutative on S . But the inversion in S is not continuous since $(U(\alpha, \varepsilon))^{-1} \not\subseteq U(\beta, \delta)$ for all $\alpha, \beta < \omega_1$ and $\varepsilon, \delta \in (0, 1)$.

Observe also that the subsemigroup of idempotents of the semigroup S can be identified with the discrete sum of $[0, \omega_1) \sqcup \{\omega_1\}$ and hence is sequentially compact, locally countable, and locally compact.

Also we observe that the projective map $\pi: S \rightarrow E(S)$ is not continuous.

Remark 15. Example 14 shows that the requirement of regularity in Theorems 2 and Corollary 2 is essential and cannot be replaced by the quasi-regularity. This contrasts with the case of paratopological groups - see [22].

ACKNOWLEDGEMENTS

This research was supported by the Slovenian Research Agency grants P1-0292-0101-04, J1-9643-0101 and BI-UA/07-08/001. The authors gratefully acknowledge the referee for several comments and suggestions which have considerably improve the original version of the manuscript.

REFERENCES

- [1] T. Banakh and S. Dimitrova, Openly factorizable spaces and compact extensions of topological semigroup, preprint (arXiv: 0811.4272).
- [2] T. Banakh, S. Dimitrova and O. Gutik, The Rees-Suschkewitsch Theorem for simple topological semigroups, *Mat. Stud.* **31**:2 (2009), in print. (preprint arXiv:0811.1371)
- [3] T. Banakh, S. Dimitrova and O. Gutik, Embedding the bicyclic semigroup into countably compact topological semigroups, preprint (arXiv: 0811.4276).

- [4] T. O. Banach and O. V. Gutik, On the continuity of inversion in countably compact inverse topological semigroups, *Semigroup Forum* **68**:3 (2004), 411—418.
- [5] B. Bokalo and I. Guran, Sequentially compact Hausdorff cancellative semigroup is a topological group, *Mat. Stud.* **6** (1996), 39—40.
- [6] N. Brand, Another note on the continuity of the inverse, *Archiv Math.* **39** (1982), 241—245.
- [7] J. H. Carruth, J. A. Hildebrandt and R. J. Koch, *The Theory of Topological Semigroups*, Vol. I., Marcell Dekker, Inc., New York and Basel, 1983; Vol. II., Marcell Dekker, Inc., New York and Basel, 1986.
- [8] A. H. Clifford and G. B. Preston, *The Algebraic Theory of Semigroups*, Vol. I., Amer. Math. Soc. Surveys 7, 1961; Vol. II. Amer. Math. Soc. Surveys 7, 1967.
- [9] C. Eberhart and J. Selden, On the closure of the bicyclic semigroup, *Trans. Amer. Math. Soc.* **144** (1969), 115—126.
- [10] R. Ellis, Continuity and homeomorphism groups, *Proc. Amer. Math. Soc.* **47** (1953), 969—977.
- [11] R. Engelking, *General Topology*, 2nd ed., Heldermann, Berlin, 1989.
- [12] S. P. Franklin, Spaces in which sequences suffice, *Fundam. Math.* **57** (1965), 107—115.
- [13] D. L. Grant, Sequentially compact cancellative topological semigroups: some progress on the Wallace problem, *Papers on General Topology and Applications (Madison, WI, 1991)*, Ann. New York Acad. Sci., 704, New York Acad. Sci., New York, 1993, pp. 150—154.
- [14] O. Gutik and D. Repovš, On countably compact 0-simple topological inverse semigroups, *Semigroup Forum* **75**:2 (2007), 464—469.
- [15] J. A. Hildebrandt and R. J. Koch, Swelling actions of Γ -compact semigroups, *Semigroup Forum* **33** (1988), 65—85.
- [16] P. S. Kenderov, I. S. Kortezov and W. B. Moors, Topological games and topological groups, *Topology Appl.* **109** (2001), 157—165.
- [17] P. Koszmider, A. Tomita and S. Watson, Forcing countably compact group topologies on a larger free Abelian group, *Topology Proceedings* **25**, Summer (2000), 563—574.
- [18] R. E. Madariaga-Garcia and A. H. Tomita, Countably compact topological group topologies on free Abelian groups from selective ultrafilters, *Topology Appl.* **154** (2007), 1470—1480.
- [19] K. Numakura, On bicomact semigroups, *Math. J. Okayama Univ.* **1** (1952), 99—108.
- [20] A. B. Paalman-de-Miranda, *Topological Semigroups*, Mathematical Centre Tracts. Vol. 11. Mathematisch Centrum, Amsterdam, 1964.
- [21] O. Ravsky, An example of a Hausdorff countably compact paratopological group which is not a topological group, Proc. IV Int. Algebraic Conf. in Ukraine, Lviv, 2003, p. 192.
- [22] O. Ravsky and E. Reznichenko, The continuity of the inverse in groups, *Intern. Conf. Func. Anal. Appl. Dedicated to 110th Anniversary of S. Banach, Lviv, Ukraine, May 28–31, 2002*, Lviv, 2002, pp. 170—172.
- [23] D. Robbie and S. Svetlichny, An answer to A. D. Wallace's question about countably compact cancellative semigroups, *Proc. Amer. Math. Soc.* **124**:1 (1996), 325—330.
- [24] A. H. Tomita, The Wallace problem: A counterexample from $\text{MA}_{\text{countable}}$ and p -compactness, *Canad. Math. Bull.* **39**:4 (1996), 486—498.
- [25] A. Suschkewitsch, Über die endlichen Gruppen, *Math. Ann.* **99** (1928), 529—547.
- [26] A. D. Wallace, The structure of topological semigroups, *Bull. Amer. Math. Soc.* **61** (1955), 95—112.
- [27] A. D. Wallace, The Suschkewitsch-Rees structure theorem for compact simple semigroups, *Proc. Nat. Acad. Sci.* **42** (1956), 430—432.
- [28] A. Weil, Sur les à Structure Uniforme et sur la Topologie Générale, *Publ. Math. Univ. Strasbourg*, Hermann, Paris, 1937.
- [29] A. A. Yur'eva, Countably compact sequential topological semigroup with two-sides cancellations is a topological group, *Mat. Stud.* **2** (1993), 23—24.

DEPARTMENT OF MECHANICS AND MATHEMATICS, IVAN FRANKO LVIV NATIONAL UNIVERSITY, UNIVERSYTETSKA 1, LVIV, 79000, UKRAINE

E-mail address: o_gutik@franko.lviv.ua, ovgutik@yahoo.com

INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, AND FACULTY OF NATURAL SCIENCES AND MATHEMATICS, UNIVERSITY OF MARIBOR, GOSPOSVETSKA 84, MARIBOR 2000, SLOVENIA

E-mail address: dusan.pagon@uni-mb.si

FACULTY OF MATHEMATICS AND PHYSICS, AND FACULTY OF EDUCATION, UNIVERSITY OF LJUBLJANA, P. O. B. 2964, LJUBLJANA, 1001, SLOVENIA

E-mail address: dusan.repovs@guest.arnes.si